

# SIDE EFFECT MANAGEMENT FOR:

**Olysio**  
(Simeprevir) 150mg capsule with food

**Sovaldi**  
(Sofosbuvier) 400mg tablet taken with or without food



## Photosensitivity:

---

- Apply a sunscreen with (UVA & UVB) protection, and an Spf block of 50 or greater. Also, wear a hat, sunglasses, and protective clothing when you will be exposed to sunlight.

## Fatigue:

---

- Drink plenty of fluids, mainly consisting of water: 2-4 liters of liquids per day, unless otherwise instructed by your provider. Staying hydrated helps to minimize fatigue.
- The bed is better saved for night-time to sleep so try, if you can, to find somewhere where you can stretch out and relax other than your bed. At least try to move your joints regularly, because even this minimum exercise is better than nothing.

## Rest and pace yourself:

---

- Rest in the early stages of the illness, but avoid complete bed rest unless absolutely necessary.
- Cautiously experiment to determine the level of activity (physical and mental) you can manage without causing fatigue.
- Keep within those limits until several days have passed, then if you're still feeling better gradually increase them.
- Write down your symptoms to see how they fluctuate in relation to your activity levels. This should help you to work out how much you can do.
- Do not be deterred by feelings of healthy tiredness (without nausea or prolonged pain), which after unaccustomed activity are normal and beneficial.
- You should however stop if you experience feelings described by some as 'flu-like' symptoms, or if prolonged pain reoccurs.
- When you feel that these feelings are well past, start your activities again at a level you can cope with and gradually build up.
- The symptoms generally fluctuate; take advantage of those days when you have more energy by doing some of the normal things you did before.

## Headache:

---

- Tylenol (acetaminophen) 2000mg max per day (4 extra strength tablets max per day)
- Apply ice packs or heat on your neck and head.
- Gently massage the muscles of your neck and scalp.
- Use relaxation exercises.
- Take aspirin, or an aspirin substitute if acetaminophen is intolerable or does not work.
- Use a decongestant medication, if you have nasal congestion.
- Reduce emotional and physical stressors, such as anger, eyestrain, or continuous loud noise.
- Avoid foods that may trigger headaches, such as aged cheeses, chocolate, nuts, red wine, alcohol, avocados, figs, raisins, or pickled foods..

## Insomnia:

---

- Chamomile Tea, Melatonin, Warm bath
- Use the bedroom only for sleeping and sex. Don't work, watch TV, or use your computer or smartphone. The goal is to associate the bedroom with sleep and sex, so that when you get in bed your brain and body get a strong signal that it's time to nod off or be intimate.
- Leave the bed when you can't sleep. Don't try to force yourself to sleep. Tossing and turning will only increase the anxiety. Get up, leave the bedroom, and do something relaxing, such as reading, drinking a warm cup of caffeine-free tea, taking a bath, or listening to soothing music. When you're sleepy, go back to bed.
- Move bedroom clocks out of view. The light from the alarm clock will affect your sleep. Watching the time go by when you can't sleep, knowing that you're going to be exhausted when the alarm goes off, will cause insomnia. You can use an alarm, but make sure you can't see the time when you're in bed.
- Make sure your bedroom is quiet, dark, and cool. Noise, light, and heat can interfere with sleep. Try using a sound machine or earplugs to hide outside noise, an open window or fan to keep the room cool, and blackout curtains or a sleep mask to block out light.
- Stick to a regular sleep schedule. Support your biological clock by going to bed and getting up at the same time every day, including weekends, even if you're tired. This will help you get back in a regular sleep rhythm.

## Insomnia (continued):

---

- Avoid naps. Napping during the day can make it more difficult to sleep at night. If you feel like you have to take a nap, limit it to 30 minutes before 3 p.m.
- Avoid stimulating activity and stressful situations before bedtime. This includes vigorous exercise; big discussions or arguments; and TV, computer, or video game use. Instead, focus on quiet, soothing activities, such as reading, knitting, or listening to soft music, while keeping lights low.
- Don't read from a backlit device (such as an iPad). If you use an e-Reader, choose one that is not backlit.
- Avoid caffeine, alcohol, and nicotine. Stop drinking caffeinated beverages at least eight hours before bed. Avoid drinking alcohol in the evening; while alcohol can make you feel sleepy, it interferes with the quality of your sleep. Quit smoking or avoid it at night, as nicotine is a stimulant.

## Nausea:

---

- Kaopectate, Pepto-Bismol, Bonine (Meclizine), Nauzene

## Diarrhea:

---

- Imodium, tucks or baby wipes (helps with irritation)
- Drinking clear fluids like water, broth, or electrolyte beverages (e.g. Pedialyte) to prevent becoming dehydrated

## Pruritus / Itching:

---

- Antihistamine such as Benadryl, or a second generation non-sedating antihistamine such as an allergy pill (Claritin, Allegra, or Zyrtec)

## Asthenia / Difficulty Breathing:

---

- You need appropriate periods of rest before 3 p.m., good hydration and nutrition and regular medical care..

## Rash:

---

- Hydrocortisone cream, Benadryl cream, Calamine lotion.

## Muscle Aches & Joint Pain / Myalgia:

---

- Tylenol (take into consideration other medication that may contain acetaminophen). Moderate exercise boosts the body's natural production of serotonin (the mood-enhancing brain chemical). It also improves blood flow to muscles, releases stress and tightness, improves sleep, all of which helps relieve pain.

## Decreased appetite:

---

- Focus on eating just one large meal per day (breakfast), with light snacks until night and have a light dinner:
- Eating frequent small meals can be helpful, and are usually easier on the stomach than large meals. To ensure you are getting enough nutrients from food, meals should be high in calories and protein. You may also want to try liquid protein drinks.

## Mouth Sores:

---

- Mix together equal amounts of Milk of Magnesia or Kaopectate and Benadryl. Milk of Magnesia and Kaopectate both contain ingredients that coat wet tissues, such as the stomach and in the mouth. Benlyn and Benadryl contain ingredients that act as mild topical anesthetics and antihistamines (which reduce inflammation). Apply the mixture to the canker sore using a cotton swab. Do not to swallow the mixture you could end up numbing the reflex that keeps the windpipe closed when you swallow.
- You can find over-the-counter antiseptic creams, lozenges and mouthwashes at your local pharmacy to help relieve canker sore pain. Examples: Orabase with Benzocaine, which covers the surface of the sore like an "oral bandage." Keeping it coated will help prevent it from getting infected.
- You can also combine 1 teaspoon salt, 1 teaspoon baking soda and 2 ounces hydrogen peroxide. Mix and rinse your mouth with it four times daily. If the taste is too strong, the tingle uncomfortable, you can dilute it with 2 ounces water. You can also just rinse your mouth with lukewarm salt water.
- Sage is another option. Used most often to spice up turkey stuffing, this herb can be used to calm an angry canker. Simply add 3 teaspoons sage leaves to 1 pint boiling water. Cover and steep for 15 minutes. Rinse your mouth with the liquid several times a day. You can also rub sage leaves into a powder and apply them directly to your sore.

## Birth Control:

---

- Oral contraceptive and barrier control method must be used during the course of therapy and six months post therapy [when taking ribavirin].

## Herbal supplements:

---

- Do not take Milk Thistle or St. John's Wart. There are no additional studies with other herbal supplements..

**Committed • Accessible • Reliable • Experienced**

HepCare Specialty Pharmacy • 2200 Park Bend Drive, Bldg. 1 • Suite 300 • Austin, TX 78758  
Office Phone: 512.381.1708 • Toll Free Number [855] 241-6658 • Fax Number: 512.490.6515  
[www.HepCarePharmacy.com](http://www.HepCarePharmacy.com)