

RHEUMATOLOGY ENROLLMENT FORM

DATE: _____ SHIP TO:
DATE NEEDED: _____ PATIENT OFFICE

PATIENT INFO
NAME _____ E-MAIL _____ DOB _____ MALE FEMALE
ADDRESS _____ CITY _____ STATE _____ ZIP _____
HOME TELEPHONE _____ MOBILE PHONE _____ SS# _____

PLEASE FAX COPY OF INSURANCE CARD (FRONT & BACK)
DIAGNOSIS CODES M06.9 Rheumatoid Arthritis M08.00 Juvenile Idiopathic Arthritis M45.9 Ankylosing Spondylitis L40.52 Psoriatic Arthritis
 Date of Diagnosis: _____ Other: _____

TREATMENT HISTORY New to this medicine Continued Treatment - If continuing treatment, has patient's condition improved or stabilized? Yes No
Patient Weight: _____ kg / lb TB/PPD Test Results? Negative Positive N/A Allergies? Latex Other: _____
Hepatitis B ruled out or being treated? Yes No N/A Concomitant Medications? Methotrexate Other: _____

PRIOR FAILED MEDICATION(S)
Medication _____ Length of Treatment _____ to _____ Medication _____ Length of Treatment _____ to _____
Reason for Discontinuing _____ Reason for Discontinuing _____

PRESCRIPTION INFORMATION	DIRECTIONS	QUANTITY	DIRECTIONS	QUANTITY
ACTEMRA® <input type="checkbox"/> 162mg/0.9ml PFS	<input type="checkbox"/> (wt < 100kg): Inject 162mg SC every other week <input type="checkbox"/> (wt > 100kg): Inject 162mg SC every week <input type="checkbox"/> _____ mg/kg SC every other week <input type="checkbox"/> Other: _____	<input type="checkbox"/> 2 PFS/Pen <input type="checkbox"/> 4 PFS/Pen Refills _____	ORENCIA® <input type="checkbox"/> 250mg/15ml Vial <input type="checkbox"/> 125mg/ml PFS <input type="checkbox"/> 125mg/ml ClickJect™ Pen	Starter: <input type="checkbox"/> Initial: Infuse _____ mg IV, then inject 125mg SC within 24 hrs Maintenance: <input type="checkbox"/> Inject 125mg SC once a week <input type="checkbox"/> Other: _____ Refills _____
CIMZIA® Starter: <input type="checkbox"/> Starter Kit 200mg PFS <input type="checkbox"/> 200mg/ml Vial & Supplies Maintenance <input type="checkbox"/> 200mg/ml PFS <input type="checkbox"/> 200mg/ml Vial & Supplies	Starter Directions: <input type="checkbox"/> Inject 400mg SC at weeks 0, 2, and 4 <input type="checkbox"/> Other: _____ Maintenance Directions: <input type="checkbox"/> Inject 200mg SC every other week <input type="checkbox"/> Inject 400mg SC every 4 weeks <input type="checkbox"/> Other: _____	<input type="checkbox"/> 1 PFS Kit (6x200 mg PFS) <input type="checkbox"/> 2 PFS <input type="checkbox"/> 4 Vials <input type="checkbox"/> 6 Vials <input type="checkbox"/> _____ Vials Refills _____	OTEZLA® <input type="checkbox"/> Starter/Titration Pack <input type="checkbox"/> 30mg Tablet	(Use Otezla START form for bridge dosage) Starter: <input type="checkbox"/> Take as directed on Starter Pack Maintenance Treatment (30mg) <input type="checkbox"/> Take 1 tablet by mouth TWICE a day <input type="checkbox"/> Take 1 tablet by mouth ONCE a day <input type="checkbox"/> Other: _____ Refills _____
COSENTYX® <input type="checkbox"/> 150mg/ml PFS <input type="checkbox"/> 150mg/ml Pen	Starter: <input type="checkbox"/> Ankylosing Spondylitis or Psoriatic Arthritis: Inject 150mg (1 pens / PFS) SC weekly at weeks 0,1,2,3 and 4, then maintenance dosing. Starter: <input type="checkbox"/> Psoriatic Arthritis with Coexistent Plaque Psoriasis: Inject 300mg (2 pens/PFS) SC weekly at weeks 0,1,2,3, and 4, then maintenance. Maintenance: <input type="checkbox"/> Ankylosing Spondylitis or Psoriatic Arthritis: Inject 150mg SC every 4 weeks. <input type="checkbox"/> Other: _____ Maintenance: <input type="checkbox"/> Psoriatic Arthritis with Coexistent Plaque Psoriasis: Inject 300mg SC every 4 weeks. <input type="checkbox"/> Other: _____	<input type="checkbox"/> 5 Pen / PFS <input type="checkbox"/> 10 Pens / PFS Refills _____ <input type="checkbox"/> 1 Pens / PFS <input type="checkbox"/> 2 Pens / PFS Refills _____	REMICADE® <input type="checkbox"/> 100mg/20ml Vial	Starter: Administer _____ mg/kg at 0,2, and 6 weeks, then maintenance dosing Maintenance: <input type="checkbox"/> Administer _____ mg/kg every _____ weeks <input type="checkbox"/> Other: _____ Refills _____
ENBREL® <input type="checkbox"/> 50mg/ml SureClick Pen® <input type="checkbox"/> 50mg/ml PFS <input type="checkbox"/> 25mg/0.5ml PFS <input type="checkbox"/> 25mg/0.5ml Vial	<input type="checkbox"/> Inject 50mg SC once a week <input type="checkbox"/> Inject 50mg SC twice a week <input type="checkbox"/> Inject 25mg SC twice a week <input type="checkbox"/> _____ 0.8mg/kg SC every week <input type="checkbox"/> Other: _____	<input type="checkbox"/> 4 <input type="checkbox"/> 8 Refills _____	RITUXAN® <input type="checkbox"/> 100mg/10ml Vial <input type="checkbox"/> 500mg/50ml Vial	Starter: <input type="checkbox"/> Administer 1000mg IV initially and in 2 weeks Maintenance: <input type="checkbox"/> Administer 1000 mg IV every _____ weeks <input type="checkbox"/> Other: _____ Refills _____
HUMIRA® <input type="checkbox"/> 40mg/0.8ml Pen <input type="checkbox"/> 40mg/0.8ml PFS	<input type="checkbox"/> Inject 40mg SC every other week <input type="checkbox"/> Inject 40mg SC once a week <input type="checkbox"/> Other: _____	<input type="checkbox"/> 2 <input type="checkbox"/> 4 Refills _____	SIMPONI® <input type="checkbox"/> 50mg/0.5ml SmartJect® (Pen) <input type="checkbox"/> 50mg/0.5ml PFS	<input type="checkbox"/> Inject 50mg SC once a month <input type="checkbox"/> Other: _____ Refills _____
HUMIRA® FOR UVEITIS Uveitis Starter <input type="checkbox"/> Psoriasis Starter Pen Kit Uveitis Maintenance Dose: <input type="checkbox"/> 40mg/0.8ml Pen <input type="checkbox"/> 40mg/0.8ml PFS	Uveitis Induction Dose: <input type="checkbox"/> Inject two 40mg Pens SC on day 1, then one 40mg Pen on day 8, then one 40mg Pen every other week Uveitis Maintenance Directions: <input type="checkbox"/> Inject one 40mg dose SC every other week <input type="checkbox"/> Other: _____	<input type="checkbox"/> 1 Starter Kit <input type="checkbox"/> 2 Pens/PFS <input type="checkbox"/> _____ Refills _____	SIMPONI® ARIA® <input type="checkbox"/> 50mg/4ml vial	Starter: <input type="checkbox"/> Infuse _____ mg (2mg/kg) IV at weeks 0 and 4, then maintenance dosing. Maintenance: <input type="checkbox"/> Infuse _____ mg (2mg/kg) IV every 8 weeks. <input type="checkbox"/> Other: _____ Refills _____
			STELARA® <input type="checkbox"/> 45mg/0.5ml PFS <input type="checkbox"/> 90mg/1ml PFS	Starter: <input type="checkbox"/> Inject 1 PFS SC on Day 1 Maintenance: <input type="checkbox"/> Inject 1 PFS 4 weeks after start of treatment, then every 12 weeks thereafter <input type="checkbox"/> Other: _____ Refills _____
			XELJANZ® <input type="checkbox"/> 5mg Tablet	<input type="checkbox"/> Take 5mg PO twice daily Refills _____
			XELJANZ XR® <input type="checkbox"/> 11mg Tablet	<input type="checkbox"/> Take 1 tablet PO once daily Refills _____
			GOUT AGENTS <input type="checkbox"/> KRSTEXXA® <input type="checkbox"/> 8mg/ml Vial	<input type="checkbox"/> Administer 8mg via iv infusion over 2 hours every 2 weeks: <input type="checkbox"/> Other: _____ Refills _____

INJECTION TRAINING Patient has received injection training Physician Office to provide injection training Pharmacy to provide injection training

PRESCRIBER INFORMATION
Prescriber's Name: _____ Contact Person: _____
Telephone: _____ Fax: _____ Email: _____
Office Address: _____ City: _____ State: _____ Zip: _____
NPI #: _____ DEA #: _____ TAX ID #: _____ Medicaid Provider #: _____
*
PRESCRIBER'S SIGNATURE _____ (DATE) _____ *IF BRAND DRUGS ARE PREFERRED, HANDWRITE "BRAND MEDICALLY NECESSARY" ABOVE
I authorize the Pharmacy noted above and its representatives to act as an agent to initiate and execute the insurance prior authorization process. © ReCapt, LP All rights Reserved