# Patient Education Information for: **ZEPATIER**®

® (elbasvir and grazoprevir)



#### What is ZEPATIER®?

- It is a fixed-dose combination tablet that is used to treat patients with hepatitis C, genotype I or genotype 4.
- There are two medications in each ZEPATIER<sup>®</sup> tablet: elbasvir and grazoprevir.

# How does ZEPATIER® work to fight the hepatitis C virus?

 It contains two direct acting agents to fight the hepatitis C virus and prevent the virus from replicating.

## What do I need to tell my doctor BEFORE taking ZEPATIER ®?

- If you have severe liver disease.
- If are you are taking nafcillin, ketoconazole, tacrolimus, modafinil, etravirine, cobicistat, or are on medication to treat your cholesterol, please notify your doctor before starting ZEPATIER®.
- \*\*Tell your doctor about all of the drugs you are taking including prescription, over-thecounter medications, natural products, and vitamins to make sure it is safe to take with ZEPATIER®
- \*\*Tell your about all your health problems including a history a liver problems, severe kidney problems, if you are on dialysis, are pregnant or plan to become pregnant, or if you are breastfeeding or planning to breastfeed.

### How is this drug best taken?

- Take this drug with or without a meal.
- Avoid missing doses to gain the most benefit from this drug to cure your hepatitis C infection.

# What are the possible side effects of taking this drug?

- The most common side effects include headache, feeling tired or weak, and upset stomach.
- Contact you doctor right away, if you have signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes
- Notify your doctor if you experience these side effects and they bother you and do not go away.
- These are not all the possible side effects of this drug. If you have further questions about side effects, call your doctor.

### How should I store ZEPATIER ®?

- Store at room temperature and in a dry place. Do not store in the bathroom.
- Keep in the original blister package until ready to take to help protect the pill from moisture.

### What do I do if I miss a dose?

- If you miss a dose, take the dose as soon as you remember but do not take two doses in the same day.
- Do not take extra doses or 2 doses at the same time of this medication.

All of the above information was adapted from the MERK Inc. package insert for ZEPATIER®.